
In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.

Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs BTI Interna 3.0 Platform **Soft Bone** Hard Bone (Mandible) In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final length. In extreme hard bone, utilize DAC (Densify After Cut) Protocol. Find protocol in IFU. Densah® Bur Densah® Bur Geometry Major Ø Minor Ø Pilot Bur I Bur 2 Bur 3 Bur 4 Pilot Bur I Bur 2 Bur 3 Bur 4 Bur 5 Bur 6 Bur 7 **Block Display Block Display** VT1525* VT1525* 2.5 **Tapered** Pilot **Pilot** (2.0)(2.0)0 - 0 - 0 - 0VT1525* VT1828 VS2228* **Tapered** 3 Pilot Pilot (2.0)(2.3)(2.5)

^{*}Denotes implant placement.


^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

^{*}Clinician must follow their implant systems recommended insertion torque guidelines.

In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.

Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs

ВТІ			Interna U	niversal																
					Sc	oft Bone			Hard Bone (Mandible)											
										In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final length. In extreme hard bone, utilize DAC (Densify After Cut) Protocol. Find protocol in IFU.										
Geometry	Major Ø	Minor Ø	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Densah® Bur Block Display	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Bur 5	Bur 6	Bur 7	Densah® Bur Block Display			
Straight	3.3		Pilot	VT1828 (2.3)	VS2228* (2.5)	_	_		Pilot	VT1828 (2.3)	VS2228* (2.5)	_	_	_	_	_	• • • • • • • • • • • • • • • • • • • •			
Straight	3.5		Pilot	VT1525 (2.0)	VS2228* (2.5)	_	_		Pilot	VT1828 (2.3)	VT2535* (3.0)	_	_	_	_	_				
Straight	3.75		Pilot	VT1828 (2.3)	VT2535* (3.0)	_	_		Pilot	VT1828 (2.3)	VT2535* (3.0)	_	_	_	_	_				
Straight	4		Pilot	VT1828 (2.3)	VT2838 (3.3)	VS3238* (3.5)	_		Pilot	VT1828 (2.3)	VT2838 (3.3)	VS3238* (3.5)	_	_	_	_				
Straight	4.25		Pilot	VT1828 (2.3)	VT2838 (3.3)	VS3238* (3.5)	_		Pilot	VT1828 (2.3)	VT2838 (3.3)	VS3238* (3.5)	_	_	_	_	• - • - • - • • • • • • • • • • • • • •			

^{*}Denotes implant placement.


^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

*Clinician must follow their implant systems recommended insertion torque guidelines.

In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.

Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs

BTI		Interna U	Interna Universal Plus																
					So	oft Bone			Hard Bone (Mandible)										
								In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final length. In extreme hard bone, utilize DAC (Densify After Cut) Protocol. Find protocol in IFU.											
Geometry	Major Ø	Minor Ø	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Densah® Bur Block Display	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Bur 5	Bur 6	Bur 7	Densah® Bur Block Display		
Tapered	4.5		Pilot	VT1525 (2.0)	VT2535 (3.0)	VT3545* (4.0)	_		Pilot	VT1828 (2.3)	VT2535 (3.0)	VT2838 (3.3)	VT3545* (4.0)	_	_	_			
Tapered	5		Pilot	VT1828 (2.3)	VT2838 (3.3)	VT3848* (4.3)	_		Pilot	VT1828 (2.3)	VT2535 (3.0)	VT2838 (3.3)	VT3545 (4.0)	VT3848 (4.3)	VS4248* (4.5)				
Tapered	5.5		Pilot	VT1525 (2.0)	VT2535 (3.0)	VT3545 (4.0)	VT4555* (5.0)		Pilot	VT1828 (2.3)	VT2535 (3.0)	VT2838 (3.3)	VT3545 (4.0)	VT3848 (4.3)	VT4555* (5.0)	_			

^{*}Denotes implant placement.


^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

*Clinician must follow their implant systems recommended insertion torque guidelines.

In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.

Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs

BTI			Interna W	Interna Wide														
					So	oft Bone			Hard Bone (Mandible)									
									In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final least									
Geometry	Major Ø	Minor Ø	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Densah® Bur Block Display	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Bur 5	Bur 6	Bur 7	Densah® Bur Block Display	
Tapered	5.5		Pilot	VT1525 (2.0)	VT2535 (3.0)	VT3545 (4.0)	VT4555* (5.0)		Pilot	VT1828 (2.3)	VT2535 (3.0)	VT2838 (3.3)	VT3545 (4.0)	VT3848 (4.3)	VT4555* (5.0)	_		
Tapered	6		Pilot	VT1828 (2.3)	VT2838 (3.3)	VT3848 (4.3)	VT4858* (5.3)		Pilot	VT1828 (2.3)	VT2535 (3.0)	VT2838 (3.3)	VT3545 (4.0)	VT3848 (4.3)	VT4858 (5.3)	VS5258* (5.5)		

^{*}Denotes implant placement.


^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

*Clinician must follow their implant systems recommended insertion torque guidelines.

In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.

Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs

BTI			Externa	Γiny																	
					Sc	oft Bone			Hard Bone (Mandible)												
											In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final length. In extreme hard bone, utilize DAC (Densify After Cut) Protocol. Find protocol in IFU.										
Geometry	Major Ø	Minor Ø	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Densah® Bur Block Display	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Bur 5	Bur 6	Bur 7	Densah® Bur Block Display				
Tapered	2.5		Pilot	VT1525* (2.0)	_	_	_		Pilot	VT1525* (2.0)	_	_	_	_	_	_	<u></u>				
Tapered	3		Pilot	VT1828* (2.3)	_	_	_	○ -○-○-○ ○ -○-○-○	Pilot	VT1828 (2.3)	VS2228* (2.5)	_	_	_	_	_					
Tapered	3.3		Pilot	VT1828 (2.3)	VS2228* (2.5)	_	_		Pilot	VT1828 (2.3)	VS2228* (2.5)	_	_	_	_	_					
Tapered	3.5		Pilot	VT1525 (2.0)	VS2228* (2.5)	_	_		Pilot	VT1828 (2.3)	VT2535* (3.0)	_		_	_	_					
Tapered	3.75		Pilot	VT1525 (2.0)	VT2535* (3.0)	_	_		Pilot	VT1828 (2.3)	VT2535* (3.0)	_	_	_	_	_					

^{*}Denotes implant placement.


^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

*Clinician must follow their implant systems recommended insertion torque guidelines.

In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.

Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs

BTI Externa Universal																		
Soft Bone								Hard Bone (Mandible)										
									In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final length. In extreme hard bone, utilize DAC (Densify After Cut) Protocol. Find protocol in IFU.									
Geometry	Major Ø	Minor Ø	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Densah® Bur Block Display	Pilot	Bur I	Bur 2	Bur 3	Bur 4	Bur 5	Bur 6	Bur 7	Densah® Bur Block Display	
Straight	3.75		Pilot	VT1525 (2.0)	VT2535* (3.0)	_	_		Pilot	VT1828 (2.3)	VT2535* (3.0)	_	_	_	_	_		
Straight	4		Pilot	VT1828 (2.3)	VT2838 (3.3)	VS3238* (3.5)	_		Pilot	VT1828 (2.3)	VT2838 (3.3)	VS3238* (3.5)	_	_	_	_		

^{*}Denotes implant placement.

^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

^{*}Clinician must follow their implant systems recommended insertion torque guidelines.

In Ridge Expansion cases, please oversize your osteotomy and make sure that the crest diameter is equal to or larger than the implant major diameter.

In Hard Bone (Mandible), after Finishing the Full Osteotomy Preparation, Use the Next Larger Size Densah Bur to the 3mm Laser-Mark Depth to make sure the Osteotomy Crestal Diameter is Equal to or Larger than the Implant Major (Crestal) Diameter.


Use Densah Burs in full-step increments for Sinus Lift cases. Example: 2.0mm, 3.0mm, 4.0mm, 5.0mm

Pilot

(2.3)

(3.3)

(4.3)





Use large block display to compare Bur identification system when using the schematic below for proper Bur usage

VT5 Set

○ VT8 Set

VS8 Set

Densifying Mode CCW (800-1500) RPMs / Cutting Mode CW (800-1500) RPMs BTI Externa Universal Plus **Soft Bone** Hard Bone (Mandible) In densifying mode make sure your osteotomy is 1.0 mm deeper than the actual implant final length. In extreme hard bone, utilize DAC (Densify After Cut) Protocol. Find protocol in IFU. Densah® Bur Densah® Bur Major Ø Minor Ø Pilot Bur I Bur 2 Bur 3 Bur 4 **Pilot** Bur I Bur 2 Bur 3 Bur 4 Bur 5 Bur 6 Bur 7 Geometry **Block Display Block Display** VT1525 VT2535 VT3545* VT1828 VT2535 VT2838 VT3545* 4.5 **Tapered** Pilot Pilot (3.0)(2.3)(3.0)(3.3)(2.0)(4.0)(4.0)0 - 0 - 0 - 00-0-0-0 VT1828 VT2838 VT3848* VT1828 VT2535 VT2838 VT3545 VT3848 VS4248*

5

Tapered

Pilot

(2.3)

(3.0)

(3.3)

(4.0)

(4.3)

(4.5)

^{*}Denotes implant placement.

^{*}Clinician experience and judgment should be used in conjunction with the Densifying Reference Guide recommendation and suggested use protocols.

^{*}Clinician must follow their implant systems recommended insertion torque guidelines.